

## Article

# **Does the Use of 4% Articaine and 1:100,000 Adrenaline, Rather Than 2% Lidocaine and 1:100,000 Adrenaline, Increase the Risk of Nerve Damage When Administered for Inferior Alveolar Nerve Blocks in Patients Undergoing Local Anaesthesia for Dental Treatment? A Mini Systematic Review of The Literature**

Stirrup, Paul and Crean, Stjohn

Available at <http://clock.uclan.ac.uk/23782/>

*Stirrup, Paul and Crean, Stjohn ORCID: 0000-0001-9336-8549 (2019) Does the Use of 4% Articaine and 1:100,000 Adrenaline, Rather Than 2% Lidocaine and 1:100,000 Adrenaline, Increase the Risk of Nerve Damage When Administered for Inferior Alveolar Nerve Blocks in Patients Undergoing Local Anaesthesia for Dental Treatment? A Mini Systematic Review of The Literature. British Dental Journal . ISSN 0007-0610*

It is advisable to refer to the publisher's version if you intend to cite from the work.

<http://dx.doi.org/10.1038/sj.bdj.2019.98>

For more information about UCLan's research in this area go to <http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including



University of  
Central Lancashire  
UCLan

**CLoK**

Central Lancashire online Knowledge

[www.clok.uclan.ac.uk](http://www.clok.uclan.ac.uk)



# **Does the Use of 4% Articaine and 1:100,000 Adrenaline, Rather Than 2% Lidocaine and 1:100,000 Adrenaline, Increase the Risk of Nerve Damage When Administered for Inferior Alveolar Nerve Blocks in Patients Undergoing Local Anaesthesia for Dental Treatment?**

## **A Mini Systematic Review of The Literature.**

Manuscript Approved (MSS-2018-270).

P Stirrup\*

Dept of Oral and Maxillofacial Surgery, Blackpool Victoria Hospital,  
Blackpool, FY3 2NR, UK

S Crean

School of Medicine in Dentistry, Allen Building,  
University of Central Lancashire  
Preston, PR1 2HE, UK.

\*Correspondence to Paul Stirrup

Email: [p.stirrup@nhs.uk](mailto:p.stirrup@nhs.uk)

## **Abstract**

This mini systematic review seeks to analyse the available literature and determine if a 4% articaine solution poses a greater risk of inferior alveolar and/or lingual nerve

damage compared to that of 2% lidocaine when administered for an inferior alveolar nerve block.

After a mini systematic review search of the published literature, seven suitable studies were identified, one double blind random controlled trial (DBRCT) and six retrospective cohort studies.

The DBRCT and 2 of the cohort studies concluded that 4% articaine poses no greater risk of nerve damage.

The remaining 4 cohort studies suggested that caution should be exhibited when using a 4% local anaesthetic solution rather than a 2% solution. However, these studies also concluded that no evidence exists to explain the reasons for their results.

The included articles present no conclusive evidence to suggest that 4% articaine causes more nerve damage than 2% lidocaine although some authors advise caution when using this agent.

All studies conclude that further quality research is required and it is therefore suggested that dental practitioners exhibit caution when choosing to use 4% articaine in an inferior alveolar nerve block until further scientific research has been performed

# Introduction

Since 1949, lidocaine has been recognised as the “gold standard” of local anaesthetic agents.<sup>1</sup> However, the desire to develop fast acting agents with a short half-life that also produce profound anaesthesia has led to the development of other alternatives. One example is articaine, initially synthesised in 1969 and used for the first time in clinical dental practice in Germany in 1976.

The reason for articaine’s popularity appears to be due to its efficacy. Numerous studies have shown that articaine produces a more profound anaesthesia than that of lidocaine.<sup>2,3,4,5,6,7,8</sup>

Lidocaine is an amide compound, based on a benzene ring structure ( $C_6H_6$ ). Articaine, in contrast, possesses a thiophene ring ( $C_4H_4S$ ), providing greater lipid solubility and an increased potency as a greater volume of an administered dose can enter the target neurons. Articaine’s lipid solubility has been quoted at over 4 times greater than that of lidocaine.<sup>9</sup> The same study confirmed that the onset of anaesthesia was achieved in 7.4 mins with articaine as opposed to 8.7 mins with lidocaine.<sup>9</sup> It has also been suggested that articaine provides a longer duration of anaesthesia due to its protein binding characteristics.<sup>10,11</sup>

With these attributes, it is perhaps not surprising that many studies have concluded that articaine is more efficient at producing profound anaesthesia than lidocaine.<sup>6,12,13,14,15</sup>

These papers include studies of both infiltration and nerve block anaesthesia. Other authors concluded that articaine has a faster onset than lidocaine<sup>11</sup>, and a meta-analysis has proved that articaine is 1.6 – 3.5 times more potent than lidocaine.<sup>2</sup>

Several studies have concluded that articaine should be recommended for use over lidocaine.<sup>2,6,12,16</sup> In 2007, Robertson et al concluded that both the speed of onset and the anaesthetic efficacy of articaine were superior to those of lidocaine when administered via a buccal infiltration technique in the posterior molar region.<sup>14</sup>

Another important attribute of a local anaesthetic agent is that of safety and this is perhaps where articaine compares less favourably. Since its introduction, several articles have been published warning of possible nerve damage when articaine is administered in an inferior alveolar nerve block (IANB).<sup>17,18</sup> These articles indicate a risk of causing temporary or permanent paraesthesia of the inferior alveolar nerve (IAN) but evidence also exists contradicting these claims.<sup>3,19,20</sup>

It appears, therefore, that the dental profession faces a dilemma. Should the more efficient agent be used to achieve faster, more profound anaesthesia or should the profession be wary of an agent that may have the potential to induce nerve damage?

A mini systematic review of the literature was performed by a single researcher with one, clearly focused question<sup>21</sup>. The results of the study will hopefully provide advice to the dental profession, ensuring the continued provision of safe and effective local anaesthesia.

# Methodology

The Scottish Intercollegiate Guidelines Network (SIGN) presents eight levels of evidence-based research. The SIGN tool was used in this study according to the criteria set out in Table 1.<sup>22</sup>

The development of the research question was aided using the PICOS method<sup>23</sup>, as described in Table 2.

Inclusion and exclusion criteria were applied to the literature search as outlined in Tables 3 and 4.

Basic search terms and medical sub headings terms were developed and detailed in Tables 5 and 6.

3 electronic databases were chosen to systematically search the available literature:

1. Medline with Full Text.
2. Dentistry and Oral Sciences Source.
3. The Cochrane Library

## Quality Assessment of Studies

To ensure that the random controlled trials included in the review were accurately assessed against the inclusion and exclusion criteria, the risk of bias tool as described in the Cochrane Handbook for Systemic Reviews of Intervention was applied.<sup>24</sup>

For the selected cohort studies, a Methodology Index for Non Randomised Studies (MINORS) was applied<sup>25</sup>, as described in Table 8.

A record sheet was developed, and each study was subsequently scored as directed by Slim and Nini et al 2003<sup>25</sup> as defined in Table 9.

## **Data Extraction**

Specifically designed data extraction forms were developed, allowing uniform data to be extracted under the following headings;

- Study design
- Study objectives
- Geographical origin of the study
- Clinical setting for the study
- Study funding
- Study participants – sex, age, numbers
- Type of anaesthetic agent used
- Study outcome – methods of recording and reporting nerve damage
- Comparison made between “expected” and “observed” outcomes
- Follow up periods
- Attrition bias
- Data analysis of outcomes



# Results

Data extraction and results of the mini systematic review are detailed in tables 10 – 18.

# Discussion

Malamed and Gagnon's study of 1325 participants enabled a statistical analysis of the results which indicated that the incidence of nerve damage was the same (1%) whether 4% articaine or 2% lidocaine was used as the LA agent. Indeed, this DBRCT concluded that articaine is a "*safe and effective*" local anaesthetic agent.<sup>19</sup>

Both studies conducted by Pogrel<sup>20,26</sup>, concluded that the incidence of nerve damage following the use of 4% articaine was in proportion to its market share.

However, 3 of the studies indicated that the use of 4% articaine elicited more adverse outcomes than would be expected when compared to the agent's market share.<sup>17,27,28</sup>

## Limitations and Characteristics of Included Studies

Several methodological inconsistencies exist throughout the included studies, making a direct comparison between the chosen articles difficult.

When performing a study comparing 2 pharmaceutical agents, a true comparison can only be achieved with the knowledge of the relative use of the 2 drugs within the studied population.

Haas and Lennon<sup>17</sup>, Gaffen and Haas<sup>28</sup> and Garisto, Gaffen et al<sup>27</sup> all used the “null hypothesis” developed by Ronald Fisher.<sup>29</sup>

However, the other included studies failed to indicate any comparison between expected and observed outcome events.

The creation of a “barb” on the tip of the needle resulting from contact with the bone, may also be a factor in the traumatic damage to both the IAN and LN. However, whether or not this event occurred during any of the IANBs included in the studies, the resultant mechanical damage would be the same for both LA solutions.

Of the 7 included papers, only one involves a DBRCT, 3 involve voluntary reporting of nerve damage and the remaining 3 articles elicit their information from patients who have been referred to a specialist centre for the specific reason that they are experiencing some degree of nerve damage. This clearly results in a considerable degree of reporting bias.

With incidences of nerve damage ranging from 1: 27,000 to 1: 785,000<sup>17,30</sup>, it is clear that this study’s outcome is extremely rare. To obtain statistically significant results in a DBRCT would require a clinical trial on a very large scale. This could explain the existence of only one such study since 1976.<sup>19</sup>

Both Hillerup and Jensen<sup>18</sup> and Garisto and Gaffen<sup>27</sup> make reference to the possibility of reporting bias in their papers and Gaffen and Haas<sup>28</sup> admit that “*reported incidence*

*numbers should be viewed cautiously*". In his 2007 paper, Pogrel<sup>26</sup> states that he estimates that his study represents approximately 10% of all cases of nerve damage in the given population per year. However, reporting bias for patients referred to a specialist centre would be the same for both LA solutions.

The only study that included a detailed physical examination of the patient was that of Hillerup and Jensen<sup>18</sup> using a "standardised test of neurosensory functions" by a single operator to determine the presence and extent of any reported nerve damage.<sup>31,32</sup> The remaining included studies merely noted the incidence of "reported" nerve damage.

Pogrel's studies<sup>20,26</sup>, using data from a specialist centre and Garisto and Gaffen's paper<sup>27</sup> all failed to accurately examine the patient, relying instead on the patient's own descriptions and a log of reported cases to AERS. Pogrel's description of the patient "examination" lacks sufficient detail to allow exclusion of detection bias.

The description of the reporting of an "electric shock" during the administration of the LA created notable discussion among the included authors. Four of the included papers noted the reporting of this phenomenon<sup>17,18,27,28</sup> and all included these reports in their results as a "nerve injury". The remaining 3 papers failed to mention this possible event.<sup>19,20,26</sup>

Interestingly, Hillerup and Jensen state that "*electric shock per se is probably of minor relevance for the aetiology of injection injuries*".<sup>18</sup> However, they then go on to question the cause of nerve injury, admitting that it is unknown as to whether the nerve is damaged via neurotoxicity or mechanically, via intra-fascicular injection.

Many authors are now advocating the use of 4% articaine in infiltration anaesthesia as an alternative to block anaesthesia due to the increased efficacy of this agent.<sup>33,34,35,36</sup>

The evidence presented in these studies indicates a clear efficacy advantage when using 4% articaine as a buccal infiltration compared to 2% lidocaine in an IANB. One author has even suggested that the IANB may now be an unnecessary procedure.<sup>37</sup>

### **Concentration of the LA agent**

Three of the chosen papers postulate that it may be the fact that, because articaine is administered in a 4% solution, it is the concentration of the LA solution rather than the actual pharmacology of the agent that causes damage to the nerve.<sup>17,27,28</sup> This suggestion would appear to be confirmed by another study on rat sciatic nerves, which concluded that significantly more neurotoxic injuries were observed following the direct injection into the nerve of a 4% articaine solution compared to that of a 2% solution.<sup>38</sup>

In a recent in-vitro study, articaine proved to be less neurotoxic than lidocaine, mepivacaine and prilocaine.<sup>39</sup> Indeed, previous studies have concluded that no scientific evidence exists to confirm the suggestion that articaine causes increased paraesthesia and, to date, no causal relationship has been exhibited between an anaesthetic agent's concentration and neurological damage.<sup>40,41</sup>

## **Implications for Clinical Research**

This mini systematic review confirms that controversy still exists over the safety of 4% articaine and 1:100,000 adrenaline as a dental local anaesthetic agent.

The authors of all the included papers admit that, due to the extremely rare occurrence of the outcome, a carefully performed, high quality DBRCT would have to involve such vast numbers of participants that, logistically, such a study would pose certain problems.

It is generally accepted that 4% articaine exhibits greater lipid solubility, faster onset and increased duration of anaesthesia, more profound anaesthesia and reduced toxicity than those of its counterpart, 2% lidocaine. With these favourable attributes, 4% articaine does indeed offer superior properties over 2% lidocaine but would a 2% articaine solution offer the same advantages?

Further research is required into the efficacy and safety of a 2% articaine solution. Indeed, a study in 2006 proved that the 4% articaine solution was not superior in its anaesthetic effect compared to 2% and 3% solutions of the same agent.<sup>42</sup>

## **Implications for General Dental Practice**

The highest level of evidence available to this study was that of Malamed and Gagnon's DBRCT in 2001.<sup>19</sup> Although spread over 27 sites in 2 countries, this trial unfortunately exhibited several potential areas of bias. It did, however, conclude that there was no evidence to suggest that 4% articaine posed a greater risk of nerve damage than 2%

lidocaine and that the use of 4% articaine in general dental practice can therefore be deemed safe and efficient.

3 further papers, not included in this study, also concluded that no conclusive evidence exists to suggest that 4% articaine poses a greater risk of nerve damage compared to other LA agents.<sup>3,10,12</sup>

## Conclusion

This mini systematic review of the literature has highlighted the fact that further research is required to determine the relative risks of using 4% articaine compared to 2% lidocaine in IANB's.

Clearly, the use of 4% articaine is becoming increasingly popular as a means of achieving successful dental anaesthesia and, if current trends continue, this agent may become the number one anaesthetic of choice in the future. This steady increase in popularity is likely to be due to the proven efficacy of this LA agent, benefiting both the patient and the operator. Indeed, the incidence of inferior alveolar nerve damage may reduce in the future as more evidence emerges to support infiltration anaesthesia.

With this in mind and, considering the contradictory evidence presented in this study, it is suggested that, until factual evidence becomes available, dental practitioners should consider all the potential risks and benefits of a particular LA agent prior to its administration.

## Declaration of Interests

The authors declare no conflicts of interest.

## References

1. Renton, T., 2013. Oral surgery: part 4. Minimising and managing nerve injuries and other complications. *British dental journal*, **215**(8), pp. 393-399.
2. Brandt, R.G., Anderson, P.F., McDonald, N.J., Sohn, W. and Peters, M.C., 2011. The pulpal anesthetic efficacy of articaine versus lidocaine in dentistry: a meta-analysis. *Journal Of The American Dental Association (1939)*, **142**(5), pp. 493-504.
3. Su, N., Li, C., Wang, H., Shen, J., Liu, W. and Kou, L., 2016. Efficacy and safety of articaine versus lidocaine for irreversible pulpitis treatment: A systematic review and meta-analysis of randomised controlled trials. *Australian Endodontic Journal: The Journal Of The Australian Society Of Endodontology Inc*, **42**(1), pp. 4-15.
4. Malamed, S.F., Gagnon, S. and Leblanc, D., 2000. Efficacy Of Articaine: A New Amide Local Anesthetic. *The Journal of the American Dental Association*, **131**(5), pp. 635-642.
5. Powell, V., 2012. Articaine is superior to lidocaine in providing pulpal anesthesia. *Journal Of The American Dental Association (1939)*, **143**(8), pp. 897-898.
6. Paxton, K. and Thome, D.E., 2010. Efficacy of Articaine Formulations: Quantitative Reviews. *Dental clinics of North America*, **54**(4), pp. 643-653.
7. Haase, A., Reader, A., Nusstein, J., Beck, M. and Drum, M., 2008. Comparing Anesthetic Efficacy of Articaine Versus Lidocaine as a Supplemental Buccal Infiltration of the Mandibular First Molar After an Inferior Alveolar Nerve Block. *The Journal of the American Dental Association*, **139**(9), pp. 1228-1235.
8. Kanaa, M.D., Whitworth, J.M. and Meechan, J.G., 2012. A Comparison of the Efficacy of 4% Articaine with 1:100,000 Epinephrine and 2% Lidocaine with 1:80,000 Epinephrine in Achieving Pulpal Anesthesia in Maxillary Teeth with Irreversible Pulpitis. *Journal of endodontics*, **38**(3), pp. 279-282.
9. Bartlett, G. and Mansoor, J., 2016. Articaine buccal infiltration vs lidocaine inferior dental block - a review of the literature. *British dental journal*, **220**(3), pp. 117-120.

10. Yapp, K.E., Hopcraft, M.S. and Parashos, P., 2011. Articaine: a review of the literature. *British dental journal*, **210**(7), pp. 323-329.
11. Costa, C.G., Tortamano, I.P., Rocha, R.G., Francischone, C.E. and Tortamano, N., 2005. Onset and duration periods of articaine and lidocaine on maxillary infiltration. *Quintessence international*, **36**(3), pp. 197-201.
12. Katyal, V., 2010. The efficacy and safety of articaine versus lignocaine in dental treatments: A meta-analysis. *Journal of dentistry*, **38**(4), pp. 307-317.
13. Srinivasan, N., Kavitha, M., Loganathan, C.S. and Padmini, G., 2009. Comparison of anesthetic efficacy of 4% articaine and 2% lidocaine for maxillary buccal infiltration in patients with irreversible pulpitis. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology & Endodontology*, **107**(1), pp. 133-136.
14. Robertson, D., Nusstein, J., Reader, A., Beck, M. and McCartney, M., 2007. The anesthetic efficacy of articaine in buccal infiltration of mandibular posterior teeth. *Journal of the American Dental Association (JADA)*, **138**(8), pp. 1104-1112.
15. Darawade, D.A., Kumar, S., Budhiraja, S., Mittal, M. and Mehta, T.N., 2014. A clinical study of efficacy of 4% articaine hydrochloride versus 2% lignocaine hydrochloride in dentistry. *Journal Of International Oral Health: JIOH*, **6**(5), pp. 81-83.
16. Balto, K., 2011. Administration of Articaine Anesthesia may Lead to Superior Profound Pulpal Anesthesia Compared with Lidocaine in Adult Patients. *Journal of Evidence Based Dental Practice*, **11**(4), pp. 183-184.
17. Haas, D.A. and Lennon, D., 1995. A 21 year retrospective study of reports of paresthesia following local anesthetic administration. *Journal (Canadian Dental Association)*, **61**(4), pp. 319.
18. Hillerup, S. and Jensen, R., 2006. Nerve injury caused by mandibular block analgesia. *International journal of oral and maxillofacial surgery*, **35**(9), pp. 878.
19. Malamed, S.F., Gagnon, S. and LeBlanc, D., 2001. Articaine hydrochloride: a study of the safety of a new amide local anesthetic. *Journal Of The American Dental Association (1939)*, **132**(2), pp. 177-185.
20. Pogrel, M.A., 2012. Permanent nerve damage from inferior alveolar nerve blocks: a current update. *Journal of the California Dental Association*, **40**(10), pp. 795-797.
21. Griffiths, P., 2002. Evidence informing practice: introducing the mini-review. *British Journal of Community Nursing*, **7**(1), pp. 38-40.



22. Petrisor, B.A., Keating, J. and Schemitsch, E., 2006. Grading the evidence: levels of evidence and grades of recommendation. *Injury*, **37**(4), pp. 321-327.
23. Makela, M. and Witt, K., 2005. How to read a paper: critical appraisal of studies for application in healthcare. *Singapore medical journal*, **46**(3), pp. 108-114.
24. Higgins, J.P.T., Altman, D.G., Gøtzsche, P.C., Jüni, P., Moher, D., Oxman, A.D., Savovic, J., Schulz, K.F., Weeks, L. and Sterne, J.A.C., 2011. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ (Clinical research ed.)*, **343**, pp. d5928-d5928.
25. Slim, K., Nini, E., Forestier, D., Kwiatkowski, F., Panis, Y. and Chipponi, J., 2003. Methodological index for non-randomized studies (minors): development and validation of a new instrument. *ANZ Journal of Surgery*, **73**(9), pp. 712-716.
26. Pogrel, M.A., 2007. Permanent nerve damage from inferior alveolar nerve blocks--an update to include articaine. *Journal of the California Dental Association*, **35**(4), pp. 271-273.
27. Garisto, G.A., Gaffen, A.S., Lawrence, H.P., Tenenbaum, H.C. and Haas, D.A., 2010. Occurrence of paresthesia after dental local anesthetic administration in the United States. *Journal of the American Dental Association (JADA)*, **141**(7), pp. 836-844.
28. Gaffen, A.S. and Haas, D.A., 2009. Retrospective review of voluntary reports of nonsurgical paresthesia in dentistry. *Journal (Canadian Dental Association)*, **75**(8), pp. 579-579.
29. Saxon, E., 2015. Defining the null hypothesis. *BMC Biology*, **13**, pp. 68-68.
30. Pogrel, M.A. and Thamby, S., 2000. Permanent Nerve Involvement Resulting: From Inferior Alveolar Nerve Blocks. *The Journal of the American Dental Association*, **131**(7), pp. 901-907.
31. Robinson, P.P., 1988. Observations on the recovery of sensation following inferior alveolar nerve injuries. *The British journal of oral & maxillofacial surgery*, **26**(3), pp. 177-189.
32. Robinson, P.P., Smith, K.G., Johnson, F.P. and Coppins, D.A., 1992. Equipment and methods for simple sensory testing. *The British journal of oral & maxillofacial surgery*, **30**(6), pp. 387-389.
33. Corbett, I.P., Kanaa, M.D., Whitworth, J.M. and Meechan, J.G., 2008. Articaine infiltration for anesthesia of mandibular first molars. *Journal of endodontics*, **34**(5), pp. 514-518.

34. Kanaa, M.D., Whitworth, J.M., Corbett, I.P. and Meechan, J.G., 2009. Articaine buccal infiltration enhances the effectiveness of lidocaine inferior alveolar nerve block. *International endodontic journal*, **42**(3), pp. 238-246.
35. Poorni, S., Veniashok, B., Senthilkumar, A.D., Indira, R. and Ramachandran, S., 2011. Anesthetic Efficacy of Four Percent Articaine for Pulpal Anesthesia by Using Inferior Alveolar Nerve Block and Buccal Infiltration Techniques in Patients with Irreversible Pulpitis: A Prospective Randomized Double-blind Clinical Trial. *Journal of endodontics*, **37**(12), pp. 1603-1607.
36. Leith, R., Lynch, K. and O'connell, A.C., 2012. Articaine use in children: a review. *European Archives Of Paediatric Dentistry: Official Journal Of The European Academy Of Paediatric Dentistry*, **13**(6), pp. 293-296.
37. Malamed, S.F., 2011. Is the mandibular nerve block passé? *Journal Of The American Dental Association (1939)*, **142 Suppl 3**, pp. 3S-7S.
38. Hillerup, S., Bakke, M., Larsen, J.O., Thomsen, C.E. and Gerds, T.E., 2011. Concentration-Dependent Neurotoxicity of Articaine: An Electrophysiological and Stereological Study of the Rat Sciatic Nerve. *Anesthesia and Analgesia*, **112**(6), pp. 1330-1338.
39. Malet, A., Faure, M., Deletage, N., Pereira, B., Haas, J. and Lambert, G., 2015. The comparative cytotoxic effects of different local anesthetics on a human neuroblastoma cell line. *Anesthesia and Analgesia*, **120**(3), pp. 589-596.
40. Malamed, S.F., 2006. Local anesthetics: dentistry's most important drugs, clinical update 2006. *Journal of the California Dental Association*, **34**(12), pp. 971-976.
41. Malamed, S.F., 2007. Articaine versus lidocaine: the author responds. *Journal of the California Dental Association*, **35**(6), pp. 383-385.
42. Hintze, A. and Paessler, L., 2006. Comparative investigations on the efficacy of articaine 4% (epinephrine 1:200,000) and articaine 2% (epinephrine 1:200,000) in local infiltration anaesthesia in dentistry--a randomised double-blind study. *Clinical oral investigations*, **10**(2), pp. 145-150.
43. Petrisor, B. and Bhandari, M., 2007. The hierarchy of evidence: Levels and grades of recommendation. *Indian Journal Of Orthopaedics*, **41**(1), pp. 11-15.
44. Aguiar, J., Chebroux, A., Martinez-Taboada, F. and Leece, E.A., 2015. Analgesic effects of maxillary and inferior alveolar nerve blocks in cats undergoing dental extractions. *Journal Of Feline Medicine And Surgery*, **17**(2), pp. 110-116.
45. Hung, P., Chang, H., Yang, P., Kuo, Y., Lan, W. and Lin, C., 2006. Comparison of the Gow-Gates mandibular block and inferior alveolar nerve block using a standardized

protocol. *Journal of the Formosan Medical Association = Taiwan yi zhi*, **105**(2), pp. 139-146.

46. Potocnik, I., Tomsic, M., Sketelj, J. and Bajrović, F.F., 2006. Articaine is More Effective than Lidocaine or Mepivacaine in Rat Sensory Nerve Conduction Block in vitro. *Journal of dental research*, **85**(2), pp. 162-166.

47. Sisk, A.L., 1986. Evaluation of the Akinosi mandibular block technique in oral surgery. *Journal Of Oral And Maxillofacial Surgery: Official Journal Of The American Association Of Oral And Maxillofacial Surgeons*, **44**(2), pp. 113-115.

48. Baroni, D.B., Franz-Montan, M., Cogo, K., Berto, L.A., Volpato, M.C., Novaes, P.D. and Groppo, F.C., 2013. Effect of articaine on mental nerve anterior portion: histological analysis in rats. *Acta Odontologica Scandinavica*, **71**(1), pp. 82-87.

49. Batista, D.S., Berto, L.A., Volpato, M.C., Ramacciato, J.C., Motta, R.H.L., Ranali, J. and Groppo, F.C., 2010. Anesthetic Efficacy of Articaine and Lidocaine for Incisive/Mental Nerve Block. *Journal of endodontics*, **36**(3), pp. 438-441.

50. Chopra, R., Jindal, G., Sachdev, V. and Sandhu, M., 2016. Double-Blind Crossover Study to Compare Pain Experience During Inferior Alveolar Nerve Block Administration Using Buffered Two Percent Lidocaine in Children. *Pediatric dentistry*, **38**(1), pp. 25-29.

51. Danielsson, K., Evers, H., Holmlund, A., Kjellman, O., Nordenram, A. and Persson, N.E., 1986. Long-acting local anaesthetics in oral surgery. Clinical evaluation of bupivacaine and etidocaine for mandibular nerve block. *International journal of oral and maxillofacial surgery*, **15**(2), pp. 119-126.

52. Rood, J.P., 1976. Inferior alveolar nerve blocks. The use of 5 per cent lignocaine. *British dental journal*, **140**(12), pp. 413-414.

53. Ahmad, Z.H., Ravikumar, H., Karale, R., Preethanath, R.S. and Sukumaran, A., 2014. Study of the anesthetic efficacy of inferior alveolar nerve block using articaine in irreversible pulpitis. *The Journal Of Contemporary Dental Practice*, **15**(1), pp. 71-74.

54. Kambalimath, D.H., Dolas, R.S., Kambalimath, H.V. and Agrawal, S.M., 2013. Efficacy of 4 % Articaine and 2 % Lidocaine: A clinical study. *Journal Of Maxillofacial And Oral Surgery*, **12**(1), pp. 3-10.

55. Moorthy, A. and Stassen, L.F.A., 2015. The occurrence of paraesthesia of the maxillary division of the trigeminal nerve after dental local anaesthetic use: a case report. *Journal of the Irish Dental Association*, **61**(1), pp. 34-35.

56. Choi, E., Seo, J., Jung, B. and Park, W., 2009. Diplopia after inferior alveolar nerve block anesthesia: report of 2 cases and literature review. *Oral surgery, oral medicine, oral pathology, oral radiology, and endodontics*, **107**(6), pp. e21-e24.

57. Al-Sandook, T. and Al-Saraj, A., 2010. Ocular complications after inferior alveolar nerve block: a case report. *Journal of the California Dental Association*, **38**(1), pp. 57-59.
58. Wyman, R.J., 2010. Nerve injury following a mandibular block: a case report. *Dentistry today*, **29**(7), pp. 14; author reply 14.
59. Pedlar, J., 2003. Prolonged paraesthesia. *British dental journal*, **195**(3), pp. 119-119.
60. Fowler, S. and Reader, A., 2013. Is a volume of 3.6 mL better than 1.8 mL for inferior alveolar nerve blocks in patients with symptomatic irreversible pulpitis? *Journal of endodontics*, **39**(8), pp. 970-972.
61. Steinkruger, G., Nusstein, J., Reader, A., Beck, M. and Weaver, J., 2006. The significance of needle bevel orientation in achieving a successful inferior alveolar nerve block. *Journal Of The American Dental Association (1939)*, **137**(12), pp. 1685-1691.

# Appendices

## Glossary of Abbreviations

AERS: Adverse Event Reporting System

DBRCT: Double Blind Random Controlled Trial

IAN: Inferior Alveolar Nerve

IANB: Inferior Alveolar Nerve Block

LA: Local Anaesthetic

LN: Lingual Nerve

MeSH: Medical Sub Headings

MINORS: Methodological Index for Non-Randomised Studies

PICOS: Population, Intervention, Comparator, Outcome, Studies

PRISMA: Preferred Reporting Items for Systematic reviews and Meta-Analyses

SIGN: Scottish Intercollegiate Guidelines Network

UCSF: University of California, San Francisco

# Tables

| Level of Evidence | Description of Evidence  |
|-------------------|--|
| 1++               | High quality meta-analysis, systematic reviews of RCT's or very low risk of bias RCT's   |
| 1+                | Well conducted meta-analysis, systematic reviews of RCT's or very low risk of bias RCT's   |
| 1-                | Meta-analysis, systematic reviews of RCT's or RCT's with a high risk of bias   |
| 2++               | High quality systematic reviews of cohort or case-control studies or high quality cohort or case-control studies with a very low risk of confounding bias or chance and a high probability that the relationship is causal |
| 2+                | Well conducted cohort or case-control studies with a low risk of confounding, bias or chance and a moderate probability that the relationship is causal  |
| 2-                | Cohort or case-control studies with a high risk of confounding, bias or chance and a significant risk that the relationship is not causal  |
| 3                 | Non-analytical studies. Case reports and case series   |
| 4                 | Expert opinion   |

**Table 1: The Hierarchy of Evidence. Adapted from the Scottish Intercollegiate Guidelines Network (SIGN).**<sup>43</sup>

| <b>PICOS</b>        | <b>Search Strategy Application</b>  |
|---------------------|---|
| <b>Population</b>   | Patients receiving IANB's with either 4% articaine hydrochloride + 1:100,000 adrenaline or patients receiving IANB's with 2% lidocaine + 1:100,000 adrenaline. Males and females. All ages  |
| <b>Intervention</b> | Studies involving the administration of an IANB with 4% articaine + 1:100,000 adrenaline  |
| <b>Comparison</b>   | Studies involving the administration of an IANB with 2% lidocaine + 1:100,000 adrenaline  |
| <b>Outcome</b>      | Post injection nerve damage indicated by prolonged temporary or permanent anaesthesia, paraesthesia or dysaesthesia in both the intervention and comparison groups.   |
| <b>Studies</b>      | Randomised controlled trials comparing 4% articaine + 1:100,000 adrenaline + 2% lidocaine + 1:100,000 adrenaline in IANB's.<br>Cohort studies investigating the use of 4% articaine + 1:100,000 adrenaline as a dental local anaesthetic agent in IANB's. |

**Table 2: PICOS parameters applied to the study.**

| <b>Inclusion Criteria</b>                     | <b>Reason for Inclusion</b>  |
|---|--|
| English language papers                       | No translation facility. Author only speaks English.                                   |
| Papers published since 1976                   | Articaine's first use in clinical dentistry  |
| Human subjects only                           | Relevant to general dental practice  |
| Male and female subjects                      | Maximum number of participants   |
| Global participation                          | Maximum number of participants   |
| Subjects of all ages                          | Maximum number of participants   |
| Articles involving IANB anaesthesia           | Specific to study question   |
| LA agents, lidocaine and articaine only       | Specific to study question   |
| Inferior alveolar and/or lingual nerve damage | Anatomical possibility of damage to either nerve during the administration of an IANB. |
| Permanent and/or temporary nerve damage       | Both indicators of nerve damage  |
| Suitable ethical approval obtained            | Ethical and moral issues relating to research  |
| Random Controlled Trials                      | Good quality evidence  |
| Cohort Studies                                | Large number of subjects   |

**Table 3: Search Inclusion Criteria**



| Exclusion Criteria  | Reason for Exclusion  |
|---|---|
| Articles describing only infiltration anaesthesia   | Administration of a nerve block is postulated as a cause of nerve damage            |
| Articles describing the use of anaesthetic agents other than articaine or lignocaine                              | Other anaesthetic agents not widely used in general dental practice                 |
| Studies investigating the use of articaine for “surgical dentistry”   | Possible surgical cause of nerve damage   |
| Studies investigating the use of articaine for removal of lower third molars and placement of mandibular implants | Both recognised causes of possible inferior alveolar and lingual nerve paraesthesia |
| “Sponsored” articles, unless a conflict of interest is declared   | Author bias   |
| Case studies  | Poor quality evidence   |
| Letters to editors  | Personal opinions   |

**Table 4: Search Exclusion Criteria**

articaine, carticaine, septanest, ultracaine, septocaine, dental anaesthesia, lignocaine, lidocaine, xylocaine, paraesthesia, paresthesia, anaesthesia, anesthesia, dysaesthesia, dysesthesia, trigeminal nerve injuries, damage, injury, inferior alveolar nerve, inferior dental nerve, mandibular nerve, lingual nerve

**Table 5. Basic Search Terms**

Articaine, dental anaesthesia, nerve injury

**Table 6. Medical Sub Headings Terms (MeSH Terms)**

| Search No. | Search Term  |
|------------|--|
|            |  |
| S1         | (MM "Carticaine")  |
| S2         | septanest  |
| S3         | articaine  |
| S4         | ultracaine   |
| S5         | septocaine   |
| S6         | (MM "Anesthesia, Dental+")                                     |
| S7         | lignocaine   |
| S8         | lidocaine  |
| S9         | xylocaine  |
| S10        | S1 OR S2 OR S3 OR S4 OR S5 OR S6                               |
| S11        | S7 OR S8 OR S9   |
| S12        | paraesthesia   |
| S13        | paresthesia  |
| S14        | anaesthesia  |
| S15        | anesthesia   |
| S16        | dysaesthesia   |
| S17        | dysesthesia  |
| S18        | (MM "Trigeminal Nerve Injuries+")                              |
| S19        | damage   |
| S20        | injury   |
| S21        | inferior alveolar nerve  |
| S22        | inferior dental nerve  |
| S23        | mandibular nerve   |
| S24        | lingual nerve  |
| S25        | S12 OR S13 OR S14 OR S15 OR S16 OR S17 OR S18 OR S19<br>OR S20 |
| S26        | S21 OR S22 OR S23 OR S24                                       |
| S27        | S10 AND S11 AND S25 AND S26                                    |

**Table 7: Search Strategy 18/11/16**

| Methodological Items for Non-Randomised Studies | Item Description   |
|---|--|
| Clearly stated aim                              | Relevant and precise study question, relating to available literature  |
| Inclusion of consecutive patients               | All eligible participants included in study  |
| Prospective collection of data                  | Data collected as per guidelines established prior to study commencement   |
| Endpoints appropriate to study aim              | Clear, quantifiable outcome addressing study question  |
| Unbiased endpoint                               | Blind assessment of endpoint   |
| Review period appropriate to aim                | Review period sufficient to allow outcome occurrence and measurement   |
| Attrition bias less than 5%                     | All patients should be reviewed  |
| Prospective calculation of study size           | Information regarding study population size necessary to achieve 95% confidence interval and level of statistical significance |
| Additional Items for use in Comparative Studies | Item Description   |
| Suitable control                                | "Gold standard" as per available information   |
| Contemporary groups                             | Groups studied during the same time period   |
| Baseline equivalent groups                      | Group criteria similar at start point  |
| Statistical analysis                            | Suitable statistics with confidence intervals or relative risk   |


**Table 8: Methodology Index for Non Randomised Studies (MINORS).<sup>25</sup>**

| Item Score | Reason                  |
|------------|-------------------------|
| 0          | Not reported            |
| 1          | Reported but inadequate |
| 2          | Reported and adequate   |

**Table 9. MINORS criteria scores.**

| Search No. | Search Term   | Dentistry & Oral Science | Medline   | Cochrane |
|------------|---|--------------------------|-----------|----------|
|            |   |                          |           |          |
| S1         | (MM "Carticaine")   | 2                        | 303       | 3        |
| S2         | septanest   | 2                        | 4         | 1        |
| S3         | articaïne   | 216                      | 398       | 3        |
| S4         | ultracaine  | 4                        | 47        | 9        |
| S5         | septocaine  | 6                        | 3         | 1        |
| S6         | (MM "Anesthesia, Dental+")  | 1,277                    | 5,827     | 9        |
| S7         | lignocaine  | 332                      | 2,405     | 11       |
| S8         | lidocaine   | 561                      | 25,426    | 47       |
| S9         | xylocaine   | 306                      | 713       | 1        |
| S10        | S1 OR S2 OR S3<br>OR S4 OR S5 OR<br>S6                                  | 1,429                    | 6,139     | 9        |
| S11        | S7 OR S8 OR S9  | 592                      | 26,463    | 55       |
| S12        | paraesthesia  | 117                      | 1,134     | 195      |
| S13        | paresthesia   | 31                       | 7,415     | 50       |
| S14        | anaesthesia   | 6,591                    | 65,803    | 1078     |
| S15        | anesthesia  | 6,591                    | 200,202   | 334      |
| S16        | dysaesthesia  | 24                       | 265       | 23       |
| S17        | dysesthesia   | 61                       | 1278      | 13       |
| S18        | (MM "Trigeminal Nerve Injuries+")                                       | 84                       | 833       | 13       |
| S19        | damage  | 3,284                    | 433,750   | 2,568    |
| S20        | injury  | 9,260                    | 549,161   | 2,570    |
| S21        | inferior alveolar nerve   | 1124                     | 2,102     | 13       |
| S22        | inferior dental nerve   | 78                       | 142       | 18       |
| S23        | mandibular nerve  | 568                      | 3,556     | 36       |
| S24        | lingual nerve   | 269                      | 1,298     | 18       |
| S25        | S12 OR S13 OR<br>S14 OR S15 OR<br>S16 OR S17 OR<br>S18 OR S19 OR<br>S20 | 18,767                   | 1,145,705 | 4,497    |
| S26        | S21 OR S22 OR<br>S23 OR S24   | 1,492                    | 5281      | 55       |
| S27        | S10 AND S11 AND<br>S25 AND S26  | 36                       | 170       | 2        |

**Table 10. Search Strategy and Results (performed on 30-12-16)**



**Table 11: PRISMA diagram indicating selection/inclusion process**

| Title and Author(s)   | Year | "SIGN"<br>Level of<br>Evidence | Type of study            |
|---|------|--------------------------------|--------------------------|
| A 21 Year Retrospective Study Of Reports Of Paresthesia Following Local Anesthetic Administration.<br>Hass and Lennon <sup>17</sup> | 1995 | 2-                             | Retrospective Cohort     |
| Retrospective Review Of Voluntary Reports Of Nonsurgical Paresthesia in Dentistry.<br>Gaffen and Haas <sup>28</sup>                 | 2009 | 2-                             | Retrospective Cohort     |
| Nerve Injury Caused By Mandibular Block Analgesia.<br>Hillerup and Jenson <sup>18</sup>   | 2006 | 2-                             | Retrospective Cohort     |
| Permanent Nerve Damage From Inferior Alveolar Nerve Blocks – An Update to Include Articaine.<br>Pogrel <sup>26</sup>                | 2007 | 2-                             | Retrospective Cohort     |
| Articaine Hydrochloride: a study of the safety of a new amide local anesthetic.<br>Malamed, Gagnon et al <sup>19</sup>              | 2001 | 1-                             | Random Controlled Trials |
| Occurrence of paresthesia after dental local anesthetic administration in the United States.<br>Garisto, Gaffen et al <sup>27</sup> | 2010 | 2-                             | Retrospective Cohort     |
| Permanent Nerve Damage From Inferior Alveolar Nerve Blocks: A Current Update.<br>Pogrel <sup>20</sup>                               | 2012 | 2-                             | Retrospective Cohort     |

**Table 12: Included Studies**

| Article(s)  | Reason for Exclusion   |
|---|--|
| Aguiar, Chebroux et al. <sup>44</sup><br>Hung, Chang et al. <sup>45</sup><br>Potocnik, Tomsic et al. <sup>46</sup><br>Sisk. <sup>47</sup><br>Baroni, Franz-Montan et al. <sup>48</sup><br>Batista, Berto et al. <sup>49</sup> | <b>Incorrect Population.</b> n = 6<br>Studies on rats and cats.<br>Studies using Cow–Gates and Akinosi IANB.<br>Studies of mental and incisive nerve blocks. |
| Chopra, Jindal et al. <sup>50</sup><br>Danielsson, Evers et al. <sup>51</sup><br>Rood. <sup>52</sup>  | <b>Incorrect Intervention.</b> n = 48<br>Studies comparing Lidocaine, etidocaine and bupivacaine.  |
| Rood. <sup>52</sup>   | <b>Incorrect Comparator.</b> n = 1<br>5% lidocaine solution used in study.   |
| Ahmad, Ravikumar et al. <sup>53</sup><br>Kambalimath, Dolas et al. <sup>54</sup><br>Moorthy, Stassen. <sup>55</sup><br>Choi, Seo et al. <sup>56</sup><br>Al-Sandook, Al-Saraj. <sup>57</sup>                                  | <b>Incorrect Outcome.</b> n = 42<br>Studies measuring articaine's efficacy only.<br>Studies detailing damage to nerves other than IAN and/or LN.             |
| Choi, Seo et al. <sup>56</sup><br>Wyman. <sup>58</sup><br>Pedlar. <sup>59</sup>   | <b>Incorrect Studies.</b> n = 8<br>Case reports and letters to editors.  |
| Fowler, Reader. <sup>60</sup><br>Steinkruger, Nusstein et al. <sup>61</sup>   | <b>Articles not answering study question.</b> n = 66<br>Studies comparing volume of anaesthetic agent and injection technique.                               |

**Table 13. Examples of excluded studies**

| Criteria                              | Haas & Lennon <sup>17</sup> | Gaffen & Haas <sup>28</sup> | Hillerup & Jenson <sup>18</sup> | Pogrel <sup>26</sup> | Malamed & Gagnon <sup>19</sup> | Garisto & Gaffen <sup>27</sup> | Pogrel <sup>20</sup> |
|---------------------------------------|-----------------------------|-----------------------------|---------------------------------|----------------------|--------------------------------|--------------------------------|----------------------|
| Clearly stated aim                    | 2                           | 2                           | 2                               | 2                    | 2                              | 2                              | 2                    |
| Inclusion of consecutive patients     | 1                           | 2                           | 2                               | 2                    | 1                              | 2                              | 2                    |
| Prospective collection of data        | 2                           | 2                           | 2                               | 2                    | 2                              | 2                              | 2                    |
| Endpoint appropriate to study         | 2                           | 2                           | 2                               | 2                    | 2                              | 2                              | 2                    |
| Unbiased assessment of endpoint       | 1                           | 1                           | 1                               | 1                    | 2                              | 1                              | 1                    |
| Appropriate follow up period          | 0                           | 1                           | 2                               | 2                    | 1                              | 1                              | 2                    |
| Loss to follow up less than 5%        | 1                           | 0                           | 0                               | 0                    | 0                              | 0                              | 0                    |
| Prospective calculation of study size | 0                           | 0                           | 0                               | 0                    | 0                              | 0                              | 0                    |
| Adequate control group                | NA                          | NA                          | NA                              | NA                   | 2                              | NA                             | NA                   |
| Contemporary groups                   | NA                          | NA                          | NA                              | NA                   | 2                              | NA                             | NA                   |
| Baseline equivalence groups           | NA                          | NA                          | NA                              | NA                   | 2                              | NA                             | NA                   |
| Adequate statistical analysis         | NA                          | NA                          | NA                              | NA                   | 1                              | NA                             | NA                   |
| Total Score                           | 9                           | 10                          | 11                              | 11                   | 17                             | 10                             | 11                   |

**Table 14: MINORS Checklist for Included Studies**



| Bias  | Malamed and Gagnon <sup>19</sup>  |
|---|---|
| Random sequence generation (selection bias)     | <b>Low risk</b> “There were no statistically significant differences in the studies between the articaine and lidocaine treatment groups with respect to age, sex, weight, race distribution or the proportion of subjects undergoing simple or complex procedures” |
| Allocation concealment (selection bias)         | <b>Unclear risk</b> . Not mentioned in methodology  |
| Blinding of outcome assessment (detection bias) | <b>Unclear risk</b> . “Randomised, double-blind...” mentioned in methodology but no other details   |
| Participant awareness (performance bias)        | <b>Unclear risk</b> . Not mentioned in methodology  |
| Incomplete outcome data (attrition bias)        | <b>High risk</b> . No mention of attrition at 24 hour and 7 day follow up interviews  |
| Sponsorship (funding bias)                      | <b>Low risk</b> . “The manufacturer of the drug products used in the three trials.....providing materials and funding.” The same company manufactures both the intervention and comparator drugs.   |
| Selective reporting (reporting bias)            | <b>Unclear risk</b> . “The vast majority of these events are related by (telephone interviews with) patients and are alleged as opposed to confirmed.”  |
| Overall risk of bias                            | <b>Unclear risk</b> .   |

**Table15: Risk of Assessment Bias (adapted from Higgins, Altman et al.<sup>24</sup>).**

| Study  | Haas & Lennon <sup>17</sup>                                | Gaffen & Haas <sup>28</sup>                                | Hillerup & Jensen <sup>18</sup>  |
|--|--|--|--|
| Study publication date                                     | April 1995   | October 2009   | May 2006   |
| Study design   | Retrospective Cohort                                       | Retrospective Cohort                                       | Retrospective Cohort   |
| Study objectives   | Prolonged paraesthesia following LA in dentistry           | Prolonged paraesthesia following LA in dentistry           | Prolonged paraesthesia following LA in dentistry   |
| Geographical origin  | Ontario, Canada  | Ontario, Canada  | Denmark  |
| Study setting  | Not Stated   | Not stated   | <i>"All dental practitioners"</i>  |
| Study funding  | Not stated   | <i>"no declared financial interests"</i>                   | Not Stated   |
| Eligible study participants                                | 143, male and female, all ages                             | 172, male and female, 11-80 years                          | 52, male and female, 24 – 81 years   |
| LA agents used   | Lidocaine, articaine, prilocaine, mepivacaine, bupivacaine | Lidocaine, articaine, prilocaine, mepivacaine, bupivacaine | Lidocaine, articaine, prilocaine, mepivacaine  |
| Outcome reporting and recording                            | Voluntary reports to PLP                                   | Voluntary reports to PLP                                   | Telephone call to GDP. Type and volume of LA used. Electric shock experienced? Written questionnaires and patient interviews |
| Comparison made between "expected" and "observed" outcomes | Yes  | Yes  | No   |
| Study period   | 21 years, 1973 - 1993                                      | 10 years, 1999 - 2008                                      | 8 years, 1997 – June 2004  |
| Attrition bias   | Not stated   | Not stated   | 30 patients (58%) lost to follow up after 12 months  |
| Data analysis of outcomes                                  | Chi – square analysis                                      | Chi – square analysis                                      | Chi – square analysis  |
| Ethical approval   | Not stated   | Stated Obtained  | Not stated   |

**Table 16a: Data Extraction**

| Study                           | Pogrel <sup>26</sup>  | Malamed, Gagnon et al <sup>19</sup>   | Garisto, Gaffen et al <sup>27</sup>                             | Pogrel <sup>20</sup>  |
|---------------------------------|---|---|---|---|
| Study publication date          | April 2007  | February 2001   | July 2010   | October 2012  |
| Study design                    | Retrospective Cohort  | 3 Double Blind Random Controlled Trials   | Retrospective Cohort  | Retrospective Cohort  |
| Study objectives                | Prolonged IAN/LN paraesthesia following LA in dentistry           | Direct comparison of efficacy and safety between 4% articaine and 2% lidocaine    | Record incidence of nerve damage after LA in dentistry          | Prolonged IAN/LN paraesthesia following LA in dentistry           |
| Geographical origin             | Maxillo Facial Dept, UCSF, USA                                    | 27 sites, 8 in the UK and 19 in the USA   | USA   | Maxillo Facial Dept, UCSF, USA                                    |
| Study setting                   | Primary and secondary dental care                                 | No stated   | Voluntary reports to FDA's AERS                                 | Primary and secondary dental care                                 |
| Study funding                   | Not stated  | <i>"Materials and funding"</i> provided by manufacturers of the LA agents         | No <i>"disclosures"</i> reported by authors                     | Not stated  |
| Eligible study participants     | 57, sex and ages not stated                                       | 1325, male and female, aged 4 – 80 years  | 226, male and female, 15 - 78 years                             | 38, sex and ages not stated                                       |
| LA agents used                  | Lidocaine, articaine, prilocaine, mepivacaine, bupivacaine        | 2% Lidocaine, 4% articaine,   | Lidocaine, articaine, prilocaine, mepivacaine, bupivacaine      | Lidocaine, articaine, prilocaine, carbocaine                      |
| Outcome reporting and recording | Examination of patient at UCSF. Details of examination not stated | Interviews and telephone calls to the patients. No further details of examination | Voluntary reports to FDA's AERS. Duration of paraesthesia noted | Examination of patient at UCSF. Details of examination not stated |

|  |                                 |                                      |  |                                 |
|--|---------------------------------|--------------------------------------|--|---------------------------------|
| Comparison made between “expected” and “observed” outcomes | Yes                             | No                                   | Yes  | Yes                             |
| Study period   | 3 years.<br>01/01/03 – 31/12/05 | Not stated                           | 11 years,<br>November 1997 – August 2008                 | 6 years,<br>01/01/06 – 31/12/11 |
| Attrition bias   | Not Stated                      | 3 patients lost to follow up (0.23%) | Not stated   | Not stated                      |
| Data analysis of outcomes                                  | Narrative                       | Narrative                            | Descriptive statistical analysis                         | Narrative                       |
| Ethical approval   | Not stated                      | Stated as obtained in UK and USA     | Stated as obtained and approved by University of Toronto | Not stated                      |

**Table 16b: Data Extraction**

| Study                               | Design                               | Number of eligible participants with outcome* | Number of participants with outcome following intervention (articaine) | Number of participants with outcome following comparison (lidocaine) | Reported Outcomes   |
|-------------------------------------|--------------------------------------|---|--|--|---|
| Haas & Lennon <sup>17</sup>         | Retrospective Cohort                 | 143*  | 50   | 5  | Paraesthesia following the injection of LA in non-surgical dentsistry |
| Gaffen & Haas <sup>28</sup>         | Retrospective Cohort                 | 172*  | 109  | 23   | Non-surgical paraesthesia   |
| Hillerup & Jensen <sup>18</sup>     | Retrospective Cohort                 | 52*   | 29   | 10   | Non-surgical IAN or LN injury following a unilateral IANB             |
| Pogrel <sup>26</sup>                | Retrospective Cohort                 | 57*   | 17   | 20   | Damage to IAN or LN following an IANB                                 |
| Malamed, Gagnon et al <sup>19</sup> | Double Blind Random Controlled Trial | 13  | 8  | 5  | <i>"numbness or tingling 4 – 8 days after the procedure"</i>          |
| Garisto, Gaffen et al <sup>27</sup> | Retrospective Cohort                 | 226*  | 116  | 11   | Oral paraesthesia following dental treatment                          |
| Pogrel <sup>20</sup>                | Retrospective Cohort                 | 38*   | 14   | 10   | Damage to IAN or LN following an IANB                                 |

**Table 17: Summary of Outcome Characteristics of Included Studies**

\* In all the included studies except Malamed, Gagnon et al, agents other than articaine and lidocaine were also studied and included in the study results. The inclusion of prilocaine, mepivacaine, bupivacaine and carbocaine explains the discrepancy between the sum of the intervention (articaine) and comparison (lidocaine) participants and that of the number of eligible participants in each study.

| Study   | Haas & Lennon <sup>17</sup> | Gaffen & Haas <sup>28</sup> | Hillerup & Jensen <sup>18</sup> |
|---|-----------------------------|-----------------------------|---------------------------------|
| Number of incidences of IAN damage with articaine           | Not reported                | Not reported                | 5                               |
| Number of incidences of LN damage with articaine            | Not reported                | Not reported                | 24                              |
| Number of incidences of IAN and/or LN damage with articaine | 50 (33.6%)                  | 109 (59.9%)                 | 29 (54%)                        |
| Number of incidences of IAN damage with lidocaine           | Not reported                | Not reported                | 3                               |
| Number of incidences of LN damage with lidocaine            | Not reported                | Not reported                | 7                               |
| Number of incidences of IAN and/or LN damage with lidocaine | 5 (3.4%)                    | 23 (12.6%)                  | 10 (19%)                        |
| Expected frequency of IAN and/or LN damage with articaine*  | 5.3                         | 26.5                        | Not reported                    |
| Observed frequency of IAN and/or LN damage with articaine   | 10                          | 42                          | Not reported                    |
| Expected frequency of IAN and/or LN damage with lidocaine*  | 3.7                         | 23.8                        | Not reported                    |
| Observed frequency of IAN and/or LN damage with lidocaine   | 0                           | 6                           | Not reported                    |

**Table 18a: Summary of Study Findings**

| Study   | Pogrel <sup>26</sup> | Malamed, Gagnon et al <sup>19</sup> | Garisto, Gaffen et al <sup>27</sup> | Pogrel <sup>20</sup> |
|---|----------------------|-------------------------------------|-------------------------------------|----------------------|
| Number of incidences of IAN damage with articaine           | Not reported         | Not reported                        | Not reported                        | Not reported         |
| Number of incidences of LN damage with articaine            | Not reported         | Not reported                        | Not reported                        | Not reported         |
| Number of incidences of IAN and/or LN damage with articaine | 17 (29.8%)           | 8 (1%)                              | 116 (51.3%)                         | 14 (37%)             |
| Number of incidences of IAN damage with lidocaine           | Not reported         | Not reported                        | Not reported                        | Not reported         |
| Number of incidences of LN damage with lidocaine            | Not reported         | Not reported                        | Not reported                        | Not reported         |
| Number of incidences of IAN and/or LN damage with lidocaine | 20 (35%)             | 5 (1%)                              | 11 (4.9%)                           | 10 (26%)             |
| Expected frequency of IAN and/or LN damage with articaine*  | Not reported         | Not reported                        | 32                                  | Not reported         |
| Observed frequency of IAN and/or LN damage with articaine   | Not reported         | Not reported                        | 116                                 | Not reported         |
| Expected frequency of IAN and/or LN damage with lidocaine*  | Not reported         | Not reported                        | 130                                 | Not reported         |
| Observed frequency of IAN and/or LN damage with lidocaine   | Not reported         | Not reported                        | 10                                  | Not reported         |

\* Expected frequencies calculated using the “null hypothesis”.<sup>29</sup>

**Table 18b: Summary of Study Findings**

